

EDUCATE YOURSELF

FACTS ABOUT HOME-BASED BASELINE TESTING

Baseline testing is used worldwide to help clinicians make concussion care treatment decisions. The best comparison of a patient's neurocognitive status after a concussion is to themselves in a 'normal' state.

Over [19 million baseline tests](#) were taken around the world.

Many communities, however, do not have the necessary resources to conduct group baseline testing. [BaselineTesting.com](#) is the solution. Individuals can take a baseline test online, at home, in about 20 minutes.



RELIABILITY

Home-based testing is just as reliable as group testing. We conducted extensive, under an external IRB research that was reviewed and approved by regulators in the United States, Canada, the European Union, and Australia. The [research](#) was designed, conducted, analyzed, and validated by leading neuropsychologists in both clinical practice and in academia.

VALIDITY

Our products are based on research, facts and [regulatory review and clearance](#). While a small number of neuropsychologists do not support baseline testing, the fact is that **home-based baseline testing has been proven to be safe and effective on 4 continents with millions of test takers.**



ACCESS TO RESULTS

Many other psychological measures are completed at home along with other tests such blood pressure, EKG, blood glucose, etc. We do not provide the home-based user with test results. We do provide home-based end users with [education](#) and access to [trained healthcare professionals](#).

